arXiv:1104.4687v1 [math.CO] 25 Apr 2011

Class two 1-planar graphs with maximum degree six or seven

Xin Zhang[*][†]

School of Mathematics, Shandong University, Jinan 250100, P. R. China

Abstract

A graph is 1-planar if it can be drawn on the plane so that each edge is crossed by at most one other edge. In this note we give examples of class two 1-planar graphs with maximum degree six or seven.

Keywords: 1-planar graph, edge coloring, class two graphs.

A graph is 1-planar if it can be drawn on the plane so that each edge is crossed by at most one other edge. This notion of 1-planar graphs was introduced by Ringel [7] while trying to simultaneously color the vertices and faces of a planar graph G such that any pair of adjacent/incident elements receive different colors. The coloring problems of 1-planar graphs have been investigated in many papers such as [1–4, 9–11].

Vizing's theorem (see page 251 of [5]) states that the edge chromatic number of every nonempty graph G is either $\Delta(G)$ or $\Delta(G) + 1$. Thus we can divide all graphs into two classes. A graph G is of Class one if $\chi'(G) = \Delta(G)$ and is of Class two if $\chi'(G) = \Delta(G) + 1$. Consequently, a major question in the area of edge colorings is that of determining to which of these two classes a given graph belongs.

For a 1-planar graph G, it is proved that G is of Class one provided $\Delta(G) \geq 10$ [10], or $\Delta(G) \geq 8$ and G contains no adjacent triangles [11], or $\Delta(G) \geq 7$ and G is triangle-free [9]. Moreover, in [11], the authors conjectured that every 1-planar graph with maximum degree at least 8 is of Class 1.

In this note we aim to construct examples of class two 1-planar graphs with maximum degree six or seven. Note that every planar graph is 1-planar graph and Vizing [8] presented examples of planar graphs of Class two with maximum degree no more than five. Therefore, we can conclude that there are 1-planar graphs of Class two with maximum degree Δ for each $\Delta \leq 7$.

At first, we need an useful lemma. Recall that $\alpha'(G)$ is the edge independent number of G and $\alpha'(G) \leq \lfloor \frac{n}{2} \rfloor$, where n is the order of G.

---

[*] Email addresses: sdu.zhang@yahoo.com.cn, xinzhang@mail.sdu.edu.cn.

[†] Supported by GIIFSDU (yzc10040) and NSFC (10971121).



Fig. 1: Class two 1-planar graphs G_1 with maximum degree 6 and G_2 with maximum degree 7.

Lemma 1. (see page 258 of [5]) If G is a graph of size m such that $m > \alpha'(G)\Delta(G)$, then G is of Class two.

Now we check that the above two 1-planar graphs are of Class two by Lemma 1. It is easy to see that the order of G_1 is 25 and the size of G_1 is 73. Thus $\alpha'(G_1) \leq 12$, which implies that $|E(G_1)| = 73 > 72 \geq \alpha'(G_1)\Delta(G_1)$. Similarly, one can also see that $|E(G_2)| = 85 > 84 = \lfloor \frac{|V(G_2)|}{2} \rfloor \Delta(G_2) \geq \alpha'(G_2)\Delta(G_2)$. So by Lemma 1, G_1 and G_2 are both of Class two.

Note that all vertices besides one 2-vertex in G_1 (resp. G_2) are of degree 6 (resp. 7). So we can respectively construct a 6-regular and a 7-regular 1-planar graph by deleting the unique 2-vertex and adding an edge between the two neighbors of it. Here we would like to point out that the resulting 7-regular 1-planar graph was presented by Fabrici and Madaras in their paper [6].

References

[1] M. O. Albertson, B. Mohar. Coloring vertices and faces of locally planar graphs. Graphs and Combinatorics **22** (2006) 289–295.

[2] O. V. Borodin. Solution of Ringel's problems on the vertex-face coloring of plane graphs and on the coloring of 1-planar graphs. Diskret. Analiz **41** (1984) 12–26.

[3] O. V. Borodin. A New Proof of the 6-Color Theorem. Journal of Graph Theory **19(4)** (1995) 507–521.

[4] O. V. Borodin, A. V. Kostochka, A. Raspaud, E. Sopena. Acyclic colouring of 1-planar graphs. Discrete Applied Mathematics **114** (2001) 29–41.

[5] G. Chartrand, P. Zhang. Chromatic Graph Theory. CRC Press, 2008.

[6] I. Fabrici, T. Madaras. The structure of 1-planar graphs. Discrete Mathematics **307** (2007) 854–865.

[7] G. Ringel. Ein Sechsfarbenproblem auf der Kugel. Abh. Math. Sem. Univ, Hamburg **29** (1965) 107–117.

[8] V. G. Vizing, Critical graphs with given chromatic class, Diskret. Analiz. **5** (1965) 9–17.

[9] X. Zhang, G. Liu, J.-L. Wu. Edge coloring of triangle-free 1-planar graphs. Journal of Shandong University (Natural Science) **45(6)** (2010) 15–17.

[10] X. Zhang, J.-L. Wu. On edge colorings of 1-planar graphs. Information Processing Letters **111(3)** (2011) 124–128.

[11] X. Zhang, G. Liu. On edge colorings of 1-planar graphs without adjacent triangles. Submitted.